

Distance domination and distance irredundance in graphs

Adriana Hansberg, Dirk Meierling and Lutz Volkmann

Lehrstuhl II für Mathematik, RWTH Aachen University, 52056 Aachen, Germany
e-mail: {hansberg,meierling,volkm}@math2.rwth-aachen.de

Abstract

A set $D \subseteq V$ of vertices is said to be a (*connected*) *distance k -dominating set* of G if the distance between each vertex $u \in V - D$ and D is at most k (and D induces a connected graph in G). The minimum cardinality of a (*connected*) distance k -dominating set in G is the (*connected*) *distance k -domination number* of G , denoted by $\gamma_k(G)$ ($\gamma_k^c(G)$, respectively). The set D is defined to be a *total k -dominating set* of G if every vertex in V is within distance k from some vertex of D other than itself. The minimum cardinality among all total k -dominating sets of G is called the *total k -domination number* of G and is denoted by $\gamma_k^t(G)$. For $x \in X \subseteq V$, if $N^k[x] - N^k[X - x] \neq \emptyset$, the vertex x is said to be *k -irredundant in X* . A set X containing only k -irredundant vertices is called *k -irredundant*. The *k -irredundance number* of G , denoted by $ir_k(G)$, is the minimum cardinality taken over all maximal k -irredundant sets of vertices of G . In this paper we establish lower bounds for the distance k -irredundance number of graphs and trees. More precisely, we prove that $\frac{5k+1}{2}ir_k(G) \geq \gamma_k^c(G) + 2k$ for each connected graph G and $(2k+1)ir_k(T) \geq \gamma_k^c(T) + 2k \geq |V| + 2k - kn_1(T)$ for each tree $T = (V, E)$ with $n_1(T)$ leaves. A class of examples shows that the latter bound is sharp. The second inequality generalizes a result of Meierling and Volkmann [9] and Cyman, Lemańska and Raczek [2] regarding γ_k and the first generalizes a result of Favaron and Kratsch [4] regarding ir_1 . Furthermore, we shall show that $\gamma_k^c(G) \leq \frac{3k+1}{2}\gamma_k^t(G) - 2k$ for each connected graph G , thereby generalizing a result of Favaron and Kratsch [4] regarding $k = 1$.

Keywords: *domination, irredundance, distance domination number, total domination number, connected domination number, distance irredundance number, tree*

2000 Mathematics Subject Classification: 05C69

1 Terminology and introduction

In this paper we consider finite, undirected, simple and connected graphs $G = (V, E)$ with vertex set V and edge set E . The number of vertices $|V|$ is called the *order* of G and is denoted by $n(G)$. For two distinct vertices u and v the *distance* $d(u, v)$ between u and v is the length of a shortest path between u and v . If X and Y are two disjoint subsets of V , then the distance between X and Y is defined as $d(X, Y) = \min \{d(x, y) \mid x \in X, y \in Y\}$. The *open k -neighborhood* $N^k(X)$ of a subset $X \subseteq V$ is the set of vertices in $V \setminus X$ of distance

at most k from X and the *closed k -neighborhood* is defined by $N^k[X] = N^k(X) \cup X$. If $X = \{v\}$ is a single vertex, then we denote the (closed) k -neighborhood of v by $N^k(v)$ ($N^k[v]$, respectively). The (closed) 1-neighborhood of a vertex v or a set X of vertices is usually denoted by $N(v)$ or $N(X)$, respectively ($N[v]$ or $N[X]$, respectively). Now let U be an arbitrary subset of V and $u \in U$. We say that v is a *private k -neighbor of u with respect to U* if $d(u, v) \leq k$ and $d(u', v) > k$ for all $u' \in U - \{u\}$, that is $v \in N^k[u] - N^k[U - \{u\}]$. The *private k -neighborhood of u with respect to U* will be denoted by $PN^k[u, U]$ ($PN^k[u]$ if $U = V$).

For a vertex $v \in V$ we define the *degree* of v as $d(v) = |N(v)|$. A vertex of degree one is called a *leaf* and the number of leaves of G will be denoted by $n_1(G)$.

A set $D \subseteq V$ of vertices is said to be a (*connected*) *distance k -dominating set* of G if the distance between each vertex $u \in V - D$ and D is at most k (and D induces a connected graph in G). The minimum cardinality of a (*connected*) distance k -dominating set in G is the (*connected*) *distance k -domination number* of G , denoted by $\gamma_k(G)$ ($\gamma_k^c(G)$, respectively). Note that the distance 1-domination number $\gamma_1(G)$ is the usual *domination number* $\gamma(G)$. A set $D \subseteq V$ of vertices is defined to be a *total k -dominating set* of G if every vertex in V is within distance k from some vertex of D other than itself. The minimum cardinality among all total k -dominating sets of G is called the *total k -domination number* of G and is denoted by $\gamma_k^t(G)$. We note that the parameters $\gamma_k^c(G)$ and $\gamma_k^t(G)$ are only defined for connected graphs and for graphs without isolated vertices, respectively.

For $x \in X \subseteq V$, if $PN^k[x] \neq \emptyset$, the vertex x is said to be *k -irredundant in X* . A set X containing only k -irredundant vertices is called *k -irredundant*. The *k -irredundance number of G* , denoted by $ir_k(G)$, is the minimum cardinality taken over all maximal k -irredundant sets of vertices of G .

In 1975, Meir and Moon [10] introduced the concept of a k -dominating set (called a ' k -covering' in [10]) in a graph, and established an upper bound for the k -domination number of a tree. More precisely, they proved that $\gamma_k(T) \leq |V(T)|/(k+1)$ for every tree T . This leads immediately to $\gamma_k(G) \leq |V(G)|/(k+1)$ for an arbitrary graph G . In 1991, Topp and Volkmann [11] gave a complete characterization of the class of graphs G that fulfill the equality $\gamma_k(G) = |V(G)|/(k+1)$.

The concept of k -irredundance was introduced by Hattingh and Henning [5] in 1995. With $k = 1$, the definition of an k -irredundant set coincides with the notion of an irredundant set, introduced by Cockayne, Hedetniemi and Miller [1] in 1978. Since then a lot of research has been done in this field and results have been presented by many authors (see [5]).

In 1991, Henning, Oellermann and Swart [8] motivated the concept of total distance domination in graphs which finds applications in many situations and structures which give rise to graphs.

For a comprehensive treatment of domination in graphs, see the monographs by Haynes, Hedetniemi and Slater [6], [7].

In this paper we establish lower bounds for the distance k -irredundance number of graphs and trees. More precisely, we prove that $\frac{5k+1}{2}ir_k(G) \geq \gamma_k^c(G) + 2k$ for each connected graph G and $(2k+1)ir_k(T) \geq \gamma_k(T) + 2k \geq |V| + 2k - kn_1(T)$ for each tree $T = (V, E)$

with $n_1(T)$ leaves. A class of examples shows that the latter bound is sharp. Since $\gamma_k(G) \geq ir_k(G)$ for each connected graph G , the latter generalizes a result of Meierling and Volkmann [9] and Cyman, Lemanska and Raczek [2] regarding γ_k and the former generalizes a result of Favaron and Kratsch [4] regarding ir_1 . In addition, we show that if G is a connected graph, then $\gamma_k^c(G) \leq (2k + 1)\gamma_k(G) - 2k$ and $\gamma_k^c(G) \leq \frac{3k-1}{2}\gamma_k^t(G) - 2k$ thereby generalizing results of Duchet and Meyniel [3] for $k = 1$ and Favaron and Kratsch [4] for $k = 1$, respectively.

2 Results

First we show the inequality $\gamma_k^c \leq (2k + 1)\gamma_k - 2k$ for connected graphs.

Theorem 2.1. *If G is a connected graph, then*

$$\gamma_k^c(G) \leq (2k + 1)\gamma_k(G) - 2k.$$

Proof. Let G be a connected graph and let D be a distance k -dominating set. Then $G[D]$ has at most $|D|$ components. Since D is a distance k -dominating set, we can connect two of these components to one component by adding at most $2k$ vertices to D . Hence, we can construct a connected k -dominating set $D' \supseteq D$ in at most $|D| - 1$ steps by adding at most $(|D| - 1)2k$ vertices to D . Consequently,

$$\gamma_k^c(G) \leq |D'| \leq |D| + (|D| - 1)2k = (2k + 1)|D| - 2k$$

and if we choose D such that $|D| = \gamma_k(G)$, the proof of this theorem is complete. \square

The results given below follow directly from Theorem 2.1.

Corollary 2.2 (Duchet & Meyniel [3] 1982). *If G is a connected graph, then*

$$\gamma^c(G) \leq 3\gamma(G) - 2.$$

Corollary 2.3 (Meierling & Volkmann [9] 2005; Cyman, Lemańska & Raczek [2] 2006). *If T is a tree with n_1 leaves, then*

$$\gamma_k(T) \geq \frac{|V(T)| - kn_1 + 2k}{2k + 1}.$$

Proof. Since $\gamma_k^c(T) \geq |V(T)| - kn_1$ for each tree T , the proposition is immediate. \square

The following lemma is a preparatory result for Theorems 2.5 and 2.7.

Lemma 2.4. *Let G be a connected graph and let I be a maximal k -irredundant set such that $ir_k(G) = |I|$. If $I_1 = \{v \in I \mid v \in PN^k[v]\}$ is the set of vertices that have no k -neighbor in I , then*

$$\gamma_k^c(G) \leq (2k + 1)ir_k(G) - 2k + (k - 1)\frac{|I - I_1|}{2}.$$

Proof. Let G be a connected graph and let $I \subseteq V$ be a maximal k -irredundant set. Let

$$I_1 := \{v \in I \mid v \in PN^k[v]\}$$

be the set of vertices in I that have no k -neighbors in I and let

$$I_2 := I - I_1$$

be the complement of I_2 in I . For each vertex $v \in I_2$ let $u_v \in PN^k[v]$ be a k -neighbor of v such that the distance between v and u_v is minimal and let

$$B := \{u_v \mid v \in I_2\}$$

be the set of these k -neighbors. Note that $|B| = |I_2|$. If w is a vertex such that $w \notin N^k[I \cup B]$, then $I \cup \{w\}$ is a k -irredundant set of G that strictly contains I , a contradiction. Hence $I \cup B$ is a k -dominating set of G .

Note that $G[I \cup B]$ has at most $|I \cup B| = |I_1| + 2|I_2|$ components. From $I \cup B$ we shall construct a connected k -dominating set $D \supseteq I \cup B$ by adding at most

$$|I_2|(k-1) + (|I_1| + \lfloor \frac{|I_2|}{2} \rfloor - 1)2k + \lfloor \frac{|I_2|}{2} \rfloor (k-1)$$

vertices to $I \cup B$.

We can connect each vertex $v \in I_2$ with its corresponding k -neighbor $u_v \in B$ by adding at most $k-1$ vertices to $I \cup B$.

Recall that each vertex $v \in I_2$ has a k -neighbor $w \neq v$ in I_2 . Therefore we need to add at most $k-1$ vertices to $I \cup B$ to connect such a pair of vertices.

By combining the two observations above, we can construct a k -dominating set $D' \supseteq I \cup B$ from $I \cup B$ with at most $|I_1| + \lfloor |I_2|/2 \rfloor$ components by adding at most $(k-1)|I_2| + (k-1)\lfloor |I_2|/2 \rfloor$ vertices to $I \cup B$. Since D' is a k -dominating set of G , these components can be joined to a connected k -dominating set D by adding at most $(|I_1| + \lfloor |I_2|/2 \rfloor - 1)2k$ vertices to D' .

All in all we have shown that there exists a connected k -dominating set D of G such that

$$\begin{aligned} |D| &\leq |I_1| + 2|I_2| + (k-1)|I_2| + (k-1) \left\lceil \frac{|I_2|}{2} \right\rceil + 2k(|I_1| + \lfloor \frac{|I_2|}{2} \rfloor - 1) \\ &\leq (2k+1)|I| - 2k + (k-1)\frac{|I_2|}{2}. \end{aligned}$$

Hence, if we choose the set I such that $|I| = ir_k(G)$, the proof of this lemma is complete. \square

Since $|I_2| \leq |I|$ for each k -irredundant set I , we derive the following theorem.

Theorem 2.5. *If G is a connected graph, then*

$$\gamma_k^c(G) \leq \frac{5k+1}{2} ir_k(G) - 2k.$$

The next result follows directly from Theorem 2.5.

Corollary 2.6 (Favaron & Kratsch [4] 1991). *If G is a connected graph, then*

$$\gamma^c(G) \leq 3ir(G) - 2.$$

For acyclic graphs Lemma 2.4 can be improved as follows.

Theorem 2.7. *If T is a tree, then*

$$\gamma_k^c(T) \leq (2k + 1)ir_k(T) - 2k.$$

Proof. Let T be a tree and let $I \subseteq V$ be a maximal k -irredundant set. Let

$$I_1 := \{v \in I \mid v \in PN^k[v]\}$$

be the set of vertices in I that have no k -neighbors in I and let

$$I_2 := I - I_1$$

be the complement of I_2 in I . For each vertex $v \in I_2$ let $u_v \in PN^k[v]$ be a k -neighbor of v such that the distance between v and u_v is minimal and let

$$B := \{u_v \mid v \in I_2\}$$

be the set of these k -neighbors. Note that $|B| = |I_2|$. If w is a vertex such that $w \notin N^k[I \cup B]$, then $I \cup \{w\}$ is a k -irredundant set of G that strictly contains I , a contradiction. Hence $I \cup B$ is a k -dominating set of G .

Note that $T[I \cup B]$ has at most $|I \cup B| = |I_1| + 2|I_2|$ components. From $I \cup B$ we shall construct a connected k -dominating set $D \supseteq I \cup B$ by adding at most

$$(2k - 1)|I_2| + 2k(|I_1| - 1)$$

vertices to $I \cup B$. To do this we need the following definitions. For each vertex $v \in I_2$ let P_v be the (unique) path between v and u_v and let x_v be the predecessor of u_v on P_v . Let $I_2 = S \cup L_1 \cup L_2$ be a partition of I_2 such that

$$S = \{v \in I_2 \mid d(v, u_v) = 1\}$$

is the set of vertices of I_2 that are connected by a ‘short’ path with u_v ,

$$L_1 = \{v \in I_2 \mid N^k(x_v) \cap I_1 \neq \emptyset\}$$

is the set of vertices of I_2 that are connected by a ‘long’ path with u_v and the vertex x_v has a k -neighbor in I_1 and

$$L_2 = I_2 - (S \cup L_1)$$

is the complement of $S \cup L_1$ in I_2 . In addition, let $L = L_1 \cup L_2$. We construct D following the procedure given below.

Step 0: Set $\mathcal{I} := I_2$, $\mathcal{S} := S$ and $\mathcal{L} := L$.

Step 1: We consider the vertices in \mathcal{S} .

Step 1.1: If there exists a vertex $v \in \mathcal{S}$ such that $d(v, w) \leq k$ for a vertex $w \in \mathcal{L}$, we can connect the vertices v , u_v , w and u_w to one component by adding at most $2(k-1)$ vertices to $I \cup B$.

Set $\mathcal{I} := \mathcal{I} - \{v, w\}$, $\mathcal{S} := \mathcal{S} - \{v\}$ and $\mathcal{L} := \mathcal{L} - \{w\}$ and repeat Step 1.1.

Step 1.2: If there exists a vertex $v \in \mathcal{S}$ such that $d(v, w) \leq k$ for a vertex $w \in \mathcal{S}$ with $v \neq w$, we can connect the vertices v , u_v , w and u_w to one component by adding at most $k-1$ vertices to $I \cup B$.

Set $\mathcal{I} := \mathcal{I} - \{v, w\}$ and $\mathcal{S} := \mathcal{S} - \{v, w\}$ and repeat Step 1.2.

Step 1.3: If there exists a vertex $v \in \mathcal{S}$ such that $d(v, w) \leq k$ for a vertex $w \in I_2 - (\mathcal{S} \cup \mathcal{L})$, we can connect the vertices v and u_v to w by adding at most $k-1$ vertices to $I \cup B$.

Set $\mathcal{I} := \mathcal{I} - \{v\}$ and $\mathcal{S} := \mathcal{S} - \{v\}$ and repeat Step 1.3.

Note that after completing Step 1 the set \mathcal{S} is empty and there are at most $|I_1| + 2|I_2| - 3(r_1 + r_2) - 2r_3$ components left, where r_i denotes the number of times Step 1.i was repeated for $i = 1, 2, 3$. Furthermore, we have added at most $(k-1)(2r_1 + r_2 + r_3)$ vertices to $I \cup B$.

Step 2: We consider the vertices in L_1 .

If there exists a vertex $v \in L_1 \cap \mathcal{L}$, let $w \in I_1$ be a k -neighbor of x_v . We can connect the vertices v , u_v and w to one component by adding at most $2(k-1)$ vertices to $I \cup B$.

Set $\mathcal{I} := \mathcal{I} - \{v\}$ and $\mathcal{L} := \mathcal{L} - \{v\}$ and repeat Step 2.

Note that after completing Step 2 we have $\mathcal{L} \subseteq L_2$ and there are at most $|I_1| + 2|I_2| - 3(r_1 + r_2) - 2r_3 - 2s$ components left, where s denotes the number of times Step 2 was repeated and the numbers r_i are defined as above. Furthermore, we have added at most $(k-1)(2r_1 + r_2 + r_3 + 2s)$ vertices to $I \cup B$.

Step 3: We consider the vertices in L_2 . Recall that for each vertex $v \in L_2$ the vertex x_v has a k -neighbor $w \in I_2$ besides v .

Let v be a vertex in $L_2 \cap \mathcal{L}$ such that x_v has a k -neighbor $w \in I_2 - \mathcal{I}$. We can connect the vertices v , u_v and w by adding at most $2(k-1)$ vertices to $I \cup B$.

Set $\mathcal{I} := \mathcal{I} - \{v\}$ and $\mathcal{L} := \mathcal{L} - \{v\}$ and repeat Step 3.

Note that after completing Step 3 the sets \mathcal{I} and \mathcal{L} are empty and there are at most $|I_1| + 2|I_2| - 3(r_1 + r_2) - 2r_3 - 2s - 2t$ components left, where t denotes the number of times Step 3 was repeated and the numbers r_i and s are defined as above. Furthermore, we have added at most $(k-1)(2r_1 + r_2 + r_3 + 2s + 2t)$ vertices to $I \cup B$.

Step 4: We connect the remaining components to one component.

Let D' be the set of vertices that consists of $I \cup B$ and all vertices added in Steps 1 to 3. Since D' is a k -dominating set of G , the remaining at most $|I_1| + 2|I_2| - 3(r_1 + r_2) - 2r_3 - 2s - 2t$ components can be connected to one component by adding at most $(|I_1| + 2|I_2| - 3(r_1 + r_2) - 2r_3 - 2s - 2t - 1)2k$ vertices to D' .

After completing Step 4 we have constructed a connected k -dominating set $D \supseteq I \cup B$ by adding at most

$$(k-1)(2r_1 + r_2 + r_3 + 2s + 2t) + (|I_1| + 2|I_2| - 3(r_1 + r_2) - 2r_3 - 2s - 2t - 1)2k$$

vertices to $I \cup B$.

We shall show now that the number of vertices we have added is less or equal than $(2k-1)|I_2| + 2k(|I_1| - 1)$. Note that $|I_2| = 2r_1 + 2r_2 + r_3 + s + t$. Then

$$\begin{aligned} & (k-1)(2r_1 + r_2 + r_3 + 2s + 2t) + (|I_1| + 2|I_2| - 3(r_1 + r_2) - 2r_3 - 2s - 2t - 1)2k \\ & \quad - (2k-1)|I_2| - 2k(|I_1| - 1) \\ & = (2k+1)|I_2| - 3k(2r_1 + 2r_2 + r_3 + s + t) - k(r_3 + s + t) \\ & \quad + (k-1)(2r_1 + r_2 + r_3 + 2s + 2t) \\ & = -(k-1)(2r_1 + 2r_2 + r_3 + s + t) - k(r_3 + s + t) + (k-1)(2r_1 + r_2 + r_3 + 2s + 2t) \\ & = -(k-1)r_2 - kr_3 - s - t \\ & \leq 0. \end{aligned}$$

If we choose $|I|$ such that $|I| = ir_k(T)$, it follows that

$$\begin{aligned} \gamma_k^c(T) & \leq |D| \leq |I_1| + 2|I_2| + 2k|I_1| + (2k-1)|I_2| - 2k \\ & = (2k+1)|I| - 2k \\ & = (2k+1)ir_k(T) - 2k \end{aligned}$$

which completes the proof of this theorem. \square

As an immediate consequence we get the following corollary.

Corollary 2.8. *If T is a tree with n_1 leaves, then*

$$ir_k(G) \geq \frac{|V(T)| - kn_1 + 2k}{2k+1}.$$

Proof. Since $\gamma_k^c(T) \geq |V(T)| - kn_1$ for each tree T , the result follows directly from Theorem 2.7. \square

Note that, since $\gamma_k(G) \geq ir_k(G)$ for each graph G , Corollary 2.8 is also a generalization of Corollary 2.3. The following theorem provides a class of examples that shows that the bound presented in Theorem 2.7 is sharp.

Theorem 2.9 (Meierling & Volkmann [9] 2005; Cyman, Lemanska & Raczek [2] 2006). *Let \mathcal{R} denote the family of trees in which the distance between each pair of distinct leaves is congruent $2k$ modulo $(2k + 1)$. If T is a tree with n_1 leaves, then*

$$\gamma_k(T) = \frac{|V(T)| - kn_1 + 2k}{2k + 1}$$

if and only if T belongs to the family \mathcal{R} .

Remark 2.10. *The graph in Figure 1 shows that the construction presented in the proof of Theorem 2.7 does not work if we allow the graph to contain cycles. It is easy to see that $I = \{v_1, v_2\}$ is an ir_2 -set of G and that $D = \{u_1, u_2, x_1, x_2, x_3\}$ is a γ_2^c -set of G . Following the construction in the proof of Theorem 2.7, we have $I_1 = \emptyset$, $I_2 = \{v_1, v_2\}$ and $B = \{u_1, u_2\}$ and consequently, $D' = I_2 \cup B \cup \{x_1, x_2, x_3\}$. But $|D'| = 7 \not\leq 6 = (2 \cdot 2 + 1)|I| - 2 \cdot 2$ and D contains none of the vertices of I .*




Figure 1.

Nevertheless, we think that the following conjecture is valid.

Conjecture 2.11. *If G is a connected graph, then*

$$\gamma_k^c(G) \leq (2k + 1)ir_k(G) - 2k.$$

Now we analyze the relation between the connected distance domination number and the total distance domination number of a graph.

Theorem 2.12. *If G is a connected graph, then*

$$\gamma_k^c(G) \leq \frac{3k + 1}{2} \gamma_k^t(G) - 2k.$$

Proof. Let G be a connected graph and let D be a total k -dominating set of G of size $\gamma_k^t(G)$. Each vertex $x \in D$ is in distance at most k of a vertex $y \in D - \{x\}$. Thus we get a dominating set of G with at most $\lfloor |D|/2 \rfloor$ components by adding at most $\lfloor |D|/2 \rfloor (k - 1)$ vertices to D . As in the proof of Lemma 2.4, the resulting components can be joined to a connected k -dominating set $|D'|$ by adding at most $(\lfloor |D|/2 \rfloor - 1)2k$ vertices. Consequently,

$$\gamma_k^c(G) \leq |D'| \leq |D| + \left\lceil \frac{|D|}{2} \right\rceil (k - 1) + \left(\left\lfloor \frac{|D|}{2} \right\rfloor - 1 \right) 2k \leq \frac{3k + 1}{2} |D| - 2k = \frac{3k + 1}{2} \gamma_k^t(G) - 2k$$

and the proof is complete. \square

For distance $k = 1$ we obtain the following result.

Corollary 2.13 (Favaron & Kratsch [4] 1991). *If G is a connected graph, then*

$$\gamma^c(G) \leq 2\gamma^t(G) - 2.$$

The following example shows that the bound presented in Theorem 2.12 is sharp.

Example 2.14. *Let P be the path on $n = (3k + 1)r$ vertices with $r \in \mathbb{N}$. Then $\gamma_k^c(P) = n - 2k$, $\gamma_k^t(P) = 2r$ and thus, $\gamma_k^c(P) = \frac{3k+1}{2}\gamma_k^t(P) - 2k$.*

References

- [1] E.J. Cockayne, S.T. Hedetniemi and D.J. Miller: Properties of hereditary hypergraphs and middle graphs, *Canad. Math. Bull.* **21** (1978), 461-468.
- [2] J. Cyman, M. Lemańska and J. Raczek: Lower bound on the distance k -domination number of a tree, *Math. Slovaca* **56** (2006), no. 2, 235-243.
- [3] P. Duchet, H. Meyniel: On Hadwiger's number and the stability number, *Ann. Discrete Math.* **13** (1982), 71-74.
- [4] O. Favaron and D. Kratsch: Ratios of domination parameters, *Advances in graph theory, Vishwa, Gulbarga* (1991), 173-182.
- [5] J.H. Hattingh and M.A. Henning: Distance irredundance in graphs, *Graph Theory, Combinatorics, and Applications*, John Wiley & Sons, Inc. 1 (1995) 529-542.
- [6] T.W. Haynes, S.T. Hedetniemi and P.J. Slater: Fundamentals of Domination in Graphs, *Marcel Dekker, New York* (1998).
- [7] T.W. Haynes, S.T. Hedetniemi and P.J. Slater: Domination in Graphs: Advanced Topics, *Marcel Dekker, New York* (1998).
- [8] M.A. Henning, O.R. Oellermann and H.C. Swart: Bounds on distance domination parameters, *J. Combin. Inform. System Sci.* **16** (1991) 11-18.
- [9] D. Meierling and L. Volkmann: A lower bound for the distance k -domination number of trees, *Result. Math.* **47** (2005), 335-339.
- [10] A. Meir and J.W. Moon: Relations between packing and covering number of a tree, *Pacific J. Math.* **61** (1975), 225-233.
- [11] J. Topp and L. Volkmann: On packing and covering numbers of graphs, *Discrete Math.* **96** (1991), 229-238.